

# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/EP05/003004

International filing date: 22 March 2005 (22.03.2005)

Document type: Certified copy of priority document

Document details: Country/Office: EP  
Number: 04007510.3  
Filing date: 29 March 2004 (29.03.2004)

Date of receipt at the International Bureau: 11 April 2005 (11.04.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



**Europäisches  
Patentamt**

**European  
Patent Office**

**Office européen  
des brevets**

**Bescheinigung**

**Certificate**

**Attestation**

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

**Patentanmeldung Nr. Patent application No. Demande de brevet n°**

04007510.3

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

**R C van Dijk**





Anmeldung Nr:  
Application no.: 04007510.3  
Demande no:

Anmeldetag:  
Date of filing: 29.03.04  
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Clariant GmbH  
Brüningstrasse 50  
65929 Frankfurt am Main  
ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:  
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.  
If no title is shown please refer to the description.  
Si aucun titre n'est indiqué se référer à la description.)

Easy-dispersible concentrate ester quat compositions

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)  
revendiquée(s)  
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/  
Classification internationale des brevets:

D06M/

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of  
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL  
PL PT RO SE SI SK TR LI



29. März 2004

Clariant GmbH

2004DE 411

Dr. OT/sch

## Description

### Easy-dispersible concentrate ester quat compositions

This invention relates to an ester quat composition that is used for production of fabric softeners at lower temperatures, when compared to conventional process.

It is well known that textiles which have been washed, either in automatic or manual washing processes, and particularly those made of cellulose fibers, present a very unpleasant touch after drying. This undesired hardened feel can be overcome by treating the garments after washing in a rinsing bath with cationic compounds which contain two long aliphatic radicals in the structure.

Based on this thought, current commercial fabric softeners available in the market are still basically composed by a dispersion of dialkylic cationic substances, most of them, dialkyldimethyl ammonium salts and diester quats of triethanolamine, diethanolamine or glycerine.

The dispersion of cationic compounds, mainly those suitable for the application in fabric softeners, is a hard task to be achieved at around room temperature, due to the poor solubility/dispersibility of these raw material in cold water.

The present invention discloses a new option for working with ester quats in the production of fabric softeners, basically consisting of a high concentrated ester quat composition dispersible in water at temperatures of about 35°C.

Due to their well-known instability towards hydrolysis reaction, ester quats are rarely formulated as concentrated products in the presence of water, especially when the resulting composition is subjected to storage before final application. However, in the present patent, all advantages of water/ester quat association, which promotes the easy

dispersibility of the final mixture, can be explored, as hydrolysis is particularly kept under control by the presence of a selected pH modifier.


In general, ester quats present problems concerning the production of viscous softeners, obliging the use of thickeners to achieve a high viscosity in the final product. A significant increase in the viscosity of softener formulations is another interesting advantage that comes from the dispersion, at lower temperatures, of the composition disclosed herein, which allows a expressive reduction or even the complete removal of thickeners from final formulations. This characteristic is especially important for some countries, more frequently in Latin America and Asia, where consumers still relate the good quality of a product to its high viscosity.

An ester quat concentrate, with 75-95% of active content and ethanol and/or glycols up to 100%, is claimed in EP 0 902 008. In that patent, a cosmetic final formulation, specifically prepared from the concentrate, is also claimed, but for being a final product, the minimum amount of water required is above the maximum tolerated in the present invention, and the addition of any other additive to keep hydrolysis under control is not mentioned. In US 5,811,385, concentrated aqueous solutions of ester quat suitable for preparing laundry detergents and cosmetics are disclosed. No organic solvents are present in these concentrated aqueous solution. US 2003 158 068 discloses fabric conditioning compositions based on ester quats, with maximum total active content of 25%. In that document, an unsaturated fatty acid is always present as a viscosity stabiliser. Final aqueous softener compositions with maximum ester quat concentration of 30% are also claimed in EP 0 669 391. The invention disclosed herein regards to a high concentrate ester quat composition, with at least 50% of active matter, indicated as raw material, for easy production of commercial fabric softeners at reduced temperatures.

The purpose of this invention is to provide concentrated ester quat compositions which contain water and thus can easily be diluted even at lower temperatures. On the same time the detrimental influence of water causing hydrolysis of the ester quat is minimized by the addition of a so-called pH-modifier.


The invention relates to an ester quat composition essentially consisting of

- a) at least 50 % by weight of an ester quat compound of the formula 1



wherein:

A is a group of the formulas  $-(\text{CH}_2)_n\text{-Q-T1}$  or



Q is selected from  $-\text{O}-\text{C}(\text{O})-$ ,  $-\text{C}(\text{O})-\text{O}-$ ;

R1 is  $(\text{CH}_2)_n\text{-Q-T2}$  or T3 or R3;

R2 is  $(\text{CH}_2)_m\text{-Q-T4}$  or T5 or R3;

R3 is H,  $\text{C}_1\text{-C}_6$  alkyl,  $\text{C}_2\text{-C}_6$  alkenyl or  $\text{C}_1\text{-C}_6$  hydroxyalkyl;

T1, T2, T3, T4, T5 are independently  $\text{C}_8\text{-C}_{22}$ -alkyl or  $\text{C}_8\text{-C}_{22}$ -alkenyl;

n and m are integers from 1 to 6; and

X is an anion, such as chloride, bromide, iodide, fluoride, sulfate, methylsulfate, hydrogensulfate, carbonate, hydrogencarbonate, phosphate, mono- and di-hydrogenphosphate, pyrophosphate, metaphosphate, nitrate, phosphonate, methylphosphonate, methandisulfonate, methylsulfonate, ethylsulfonate, and other anions which are softener-compatible.

- b) an organic solvent,  
c) water and  
d) a pH modifier.



Preferred ester quat compounds of formula 1 are those wherein:

Q is  $-O-C(O)-$ ;

$R^1$  is  $(CH_2)_n-Q-T^2$  or  $T^3$ ;

$R^2$  is  $(CH_2)_m-Q-T^4$  or  $T^5$  or  $R^3$ ;

$R^3$  is  $C_1-C_6$ -alkyl, especially methyl,  $C_2-C_6$ -alkenyl or  $C_1-C_6$ -hydroxyalkyl, especially hydroxyethyl;

$T^1, T^2, T^3, T^4, T^5$  are independently  $C_8-C_{22}$ -alkyl or  $C_8-C_{22}$ -alkenyl;

n and m are 1 or 2.

Most preferred ester quats compounds are triethanolamine-diester quats and diethanolamine-diester quats that means compounds of the formula 1 wherein A,  $R^1$  and  $R^2$  are a group of the formula  $-CH_2CH_2OCO-T^1$ ,  $R^2$  in addition may be a group  $R^3$ ,  $R^2$  and  $R^3$  being independently selected  $R^3$  and X being as defined under formula 1. All mentioned ester quat compounds may contain any kind of anion, which is compatible with the ester-quat, the preferred ones are chloride, bromide, acetate, lactate, sulfate, hydrogensulfate or methylsulfate.

The claimed composition may contain these cationic compounds in an amount of from 50 to 90%, more precisely from 65 to 75% by weight of the whole composition.

b) an organic solvent

In principle, suitable organic solvents are any mono- or polyhydric short alcohols. Preference is given to using alcohols having from 1 to 4 carbon atoms, such as methanol, ethanol, propanol, isopropanol, straight chain and branched butanol, glycerol and mixtures of said alcohols. Other preferred solvents are polyethylene glycols having a relative molecular mass below 2000. In particular, the use of polyethylene glycol having a relative molecular mass between 200 and 600 and in amounts up to 30 % by weight, and of polyethylene glycol having a relative molecular mass between 400 and 600 in amounts from 5 to 30 % by weight is preferred. Also the lower alkyl ether of ethyleneglycol, propyleneglycol, polyethyleneglycol and polypropyleneglycol can be used. Other suitable solvents are, for example, triacetin (glycerol triacetate), 1-methoxy-2-propanol, hexyleneglycol.

The claimed composition may contain these organic solvents or even mixture thereof in an amount of from 5 to 30%, more precisely from 15 to 25% by weight of the whole composition.

c) water

In the present invention, its presence in the concentrate is essential to the good dispersibility at low temperatures. Accordingly, water is present in an amount of from 5 to 20%, more precisely from 7 to 15% by weight of the whole composition.

d) a pH modifier

When using ester quats as cationic compounds, the presence of water in the concentrate is a real problem, as the ester bonds are broken through the hydrolysis process. The best way to avoid it is using a controlled dosage of a pH modifier to increase the pH and thus slowing the hydrolysis significantly.

The pH modifiers especially indicated to the present composition are amine compounds, especially those selected from the group of triethanolamine, monoethanolamine, ethylenediamine, dialkylamines, dialkyl methyl amines, ethoxylated alkyl amines and aminomethyl propanol.

The claimed composition may contain a pH modifier in an amount of from 0.1 to 3% by weight of the whole composition.

Depending on the intended use, the compositions according to the invention comprise, in addition to the mentioned compounds, additives and auxiliaries which are customary and specific in each case such as for example electrolytes, perfumes, perfume carriers, colorants, hydrotropes, antifoaming agents, polymeric or other thickening agents, opacifiers, and anti-corrosion agents.

The compositions according to the invention are notable for a good storage stability and for allowing a fabric softener production by dilution with water at around room temperature.

The compositions according to the present invention can be made by melting the ester quat compound and adding the organic solvent to the monter est quat.

Water and pH-modifier are mixed separately and this mixture is added to the mixture of the est quat and the organic solvent, which mixture has been cooled down to approximately 40 to 50°C.

## Examples

### 1) Hydrolytic Stability of Easy-Dispersible Concentrate Ester Quat Compositions

The table below shows the benefits from the presence of a selected pH modifier on the hydrolysis of the Ester Quat Concentrate. The test was conducted at 35°C, which is one of the indicated temperatures for the dispersion of the product.

Percentual Acid Value Increase during Storage Test at 35°C

	after 1 week	after 2 weeks	after 3 weeks	after 4 weeks
Ester Quat Concentrate without pH modifier	31.1%	39.0%	48.0%	60.7%
Ester Quat Concentrate with pH modifier	1.3%	1.3%	3.4%	13.9%

The ester quat used in this test was C<sub>16</sub>-C<sub>18</sub>-Dialkenoyloxyethyl-hydroxyethyl-methyl-ammonium methylsulfate in isopropanol (90 % a.m.). The pH modifier was triethanolamine (99 %).

From the table, it is clear that the pH modifier plays a definitive role in keeping the hydrolysis under control during storage, mainly at higher temperatures.

### 2) Concentrate Composition (% by weight)

A) 80.00 C<sub>16</sub>-C<sub>18</sub>-Dialkanoyloxyethyl-hydroxyethyl-methyl-ammonium methylsulphate in isopropanol (90 % a.m.) (Praepagen® TQ)

- B) 10.00 Isopropanol (99.5%)
- C) 0.72 Triethanolamine (99%)
- D) 9.28 Water

Procedure:

- I. Heated A in order to get a molten material (approx. 50°C), under stirring.
- II. Added B, under stirring, and cooled to approximately 40°C (mixture 1).
- III. Mixed, in a separate recipient, C and D at room temperature (mixture 2).
- IV. Added mixture 2 into mixture 1, under stirring at 40°C.
- V. Stirred for approx. 5 minutes.

3) Concentrate Composition (% by weight)

- A) 80.00 C<sub>16</sub>-C<sub>18</sub>-Dialkanoyloxyethyl-hydroxyethyl-methyl-ammonium  
methysulphate in ethanol (85 % a.m.) (Praepagen® TQL-E)
- B) 10.00 Ethanol
- C) 0.68 Triethanolamine (99%)
- D) 9.32 Water

Procedure:

- I. Heated A in order to get a molten material (approx. 60°C), under stirring.
- II. Added B, under stirring, and cooled to approximately 50°C (mixture 1).
- III. Mixed, in a separate recipient, C and D at room temperature (mixture 2).
- IV. Added mixture 2 into mixture 1, under stirring at 50°C.
- V. Stirred for approx. 5 minutes.

4) Concentrate Composition (% by weight)

- A) 0.00 C<sub>16</sub>-C<sub>18</sub>-Dialkanoyloxyethyl-hydroxyethyl-methyl-ammonium  
thylsulphate in isopropanol (85 % a.m.) (Praepagen® TQL)
- B) 0.00 Isopropanol
- C) 0.68 Triethanolamine (99%)
- D) 9.32 Water

Procedure:

- I. Heated A in order to get a molten material (approx. 60°C), under stirring.
- II. Added, under stirring, and cooled to approximately 50°C (mixture 1).
- III. Mixed in a separate recipient, C and D at room temperature (mixture 2).
- IV. Added mixture 2 into mixture 1, under stirring at 50°C.
- V. Stirred approx. 5 minutes.

5) Concentrate Composition (% by weight)

- A) 80.00 C<sub>16</sub>-C<sub>18</sub>-Dialkanoyloxyethyl-hydroxyethyl-methyl-ammonium  
methysulphate in isopropanol (90 % a.m.) (Praepagen® TQ)
- B) 15.00 Isopropanol
- C) 0.72 Triethanolamine (99%)
- D) 4.32 Water

Procedure:

- I. Heated A in order to get a molten material (approx. 50°C), under stirring.
- II. Added B, under stirring, and cooled to approximately 40°C (mixture 1).
- III. Mixed, in a separate recipient, C and D at room temperature (mixture 2).
- IV. Added mixture 2 into mixture 1, under stirring at 40°C.
- V. Stirred for approx. 5 minutes.

6) Concentrate Composition (% by weight)

- A) 80.00 C<sub>16</sub>-C<sub>18</sub>-Dialkanoyloxyethyl-hydroxyethyl-methyl-ammonium  
methysulphate in isopropanol (90 % a.m.) (Praepagen® TQ)
- B) 15.00 Isopropanol
- C) 0.72 Aminomethyl propanol
- D) 4.32 Water

Procedure:

- I. Heated A in order to get a molten material (approx. 50°C), under stirring.
- II. Added B, under stirring, and cooled to approximately 40°C (mixture 1).
- III. Mixed, in a separate recipient, C and D at room temperature (mixture 2).

- IV. Added mixture 2 into mixture 1, under stirring at 40°C.
- V. Stirred for approx. 5 minutes.

7) Concentrate Composition (% by weight)

- A) 80.00 C<sub>16</sub>-C<sub>18</sub>-Dialkanoyloxyethyl-hydroxyethyl-methyl-ammonium methylsulphate in ethanol (90 % a.m.) (Praepagen® TQ-E)
- B) 10.00 Ethanol (99.5%)
- C) 0.72 Triethanolamine (99%)
- D) 9.28 Water

Procedure:

- I. Heated A in order to get a molten material (approx. 50°C), under stirring.
- II. Added B, under stirring, and cool to approximately 40°C (mixture 1).
- III. Mixed, in a separate recipient, C and D at room temperature (mixture 2).
- IV. Added mixture 2 into mixture 1, under stirring at 40°C.
- V. Stirred for approx. 5 minutes.

8) Softener formulations

The table below shows some comparative viscosity results between softeners prepared with an ester quat (Praepagen TQ), as normally available in the market, and with the Ester Quat Concentrate according to the present invention.

	3.5% a.m. without thickener	3.5% a.m. with 1.0% thickener	7.0% a.m. without thickener
Ester Quat (Praepagen TQ) (Procedure 1)	5	215	105
Ester Quat Concentrate (Procedure 2)	230	685	830

Viscosity (cP) – Brookfield DV-II, 20 rpm, 25°C, spindle 3

The thickener used in the example above was a modified corn starch.

Procedures:

Procedure 1 – Hot Process using conventional ester quat (Praepagen TQ):

- A) Molten the ester quat (Praepagen TQ) at approx. 65°C.
- B) Mixed the water and thickener (when present) and heated to approx. 70°C.
- C) Added the ester quat (Praepagen TQ) to the water (or mixture with thickener), under stirring.
- D) Stirred for 45 minutes under cooling.

Procedure 2 – Process with Reduced Temperatures

- A) Molten the Ester Quat Concentrate at approx. 35°C.
- B) Mixed the water and thickener (when present), at room temperature.
- C) Added the Ester Quat Concentrate to the water (or mixture with thickener), under stirring.
- D) Stirred for 30 minutes.


It is evident from the above given viscosity datas that an ester quat concentrate according to the present invention significantly increases the viscosity of the final softener composition as compared with a conventional ester quat.

## Claims

EPO - Munich  
17


29. März 2004

1. An ester quat composition essentially consisting of:  
 a) at least 50% by weight of an ester quat compound of the formula 1



wherein:

A is a group of the formulas  $-(\text{CH}_2)_n\text{-Q-T}^1$  or



Q is  $-\text{O}-\text{C}(\text{O})-$  or  $-\text{C}(\text{O})-\text{O}-$ ;

R<sup>1</sup> is  $(\text{CH}_2)_n\text{-Q-T}^2$  or T<sup>3</sup> or R<sup>3</sup>;

R<sup>2</sup> is  $(\text{CH}_2)_m\text{-Q-T}^4$  or T<sup>5</sup> or R<sup>3</sup>;

R<sup>3</sup> is H, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl or C<sub>1</sub>-C<sub>6</sub>-hydroxyalkyl;

T<sup>1</sup>, T<sup>2</sup>, T<sup>3</sup>, T<sup>4</sup>, T<sup>5</sup> are independently C<sub>8</sub>-C<sub>22</sub>-alkyl or C<sub>8</sub>-C<sub>22</sub>-alkenyl;

n and m are integers from 1 to 6; and

X is an anion,

- b) an organic solvent  
 c) water  
 d) a pH modifier.



2. A concentrated ester quat composition as claimed in claim 1, wherein in the formula 1

Q is  $-O-C(O)-$ ;

R1 is  $(CH_2)_n-Q-T2$  or T3;

R2 is  $(CH_2)_m-Q-T4$  or T5 or R3;

R3 is  $C_1-C_2$ -alkyl or  $C_1-C_6$ -hydroxyalkyl;

T1, T2, T3, T4, T5 are independently  $C_8-C_{22}$ -alkyl or  $C_8-C_{22}$ -alkenyl;

n and m are 1 or 2.

3. A concentrated ester quat composition as claimed in any of the previous claims, wherein two or more different types of the ester quat compounds of formula 1 as defined in claim 1 are present.

4. A concentrated ester quat composition as claimed in any of the previous claims, wherein the ester quat compound is a compound of formula 1 wherein A  $R^1$  and  $R^2$  are a group of the formula  $-CH_2CH_2OCO-T^1$ ,  $R^2$  in addition may be a group  $R^3$ ,  $R^2$  and  $R^3$  being independently selected,  $R^3$  and X being as defined under formula 1.

5. A concentrated ester quat composition as claimed in any of the previous claims, wherein the ester quat compound is present in an amount of from 50 to 90%, preferably from 65 to 75% by weight of the whole composition.

6. A concentrated ester quat composition as claimed in any of the previous claims, wherein the organic solvent is selected from the group of short alcohols, preferably methanol, ethanol, propanol, isopropanol or a mixture thereof.

7. A concentrated ester quat composition as claimed in any of the previous claims, wherein the organic solvent, or a mixture of organic solvents, is present in an amount of from 5 to 30%, preferably from 15 to 25% by weight of the whole composition.

8. A concentrated ester quat composition as claimed in any of the previous claims, wherein the water is present in an amount of from 5 to 20%, preferably from 7 to 15% by weight of the whole composition.
9. A concentrated ester quat composition as claimed in any one of the previous claims, wherein the pH-modifier is an amine.
10. A concentrated ester quat composition as claimed in any of the previous claims, wherein the pH modifier is selected from the group of triethanolamine, monoethanolamine, ethylenediamine, dialkylamines, dialkyl methyl amines, ethoxylated alkyl amines and methyl propanol alkyl amine.
11. A concentrated ester quat composition as claimed in any of the previous claims, wherein the pH modifier is present in an amount of from 0.1 to 3% by weight of the whole composition.
12. A concentrated ester quat composition as claimed in any of the previous claims, which additionally contains any other additives or auxiliaries.



## Summary

EPO - Munich  
17

29. März 2004

## Easy-dispersible concentrate ester quat compositions

This invention relates to an ester quat concentrate suitable for production of fabric softeners at lower temperatures comprising a) an esterquat compound; b) an organic solvent; c) water; d) a pH modifier. Although the presence of water in an ester quat concentrate is generally avoided due to the latter well known instability towards hydrolysis reaction, in the present invention, water is essential to the dispersibility at lower temperatures, which consequently leads to a significative increase in the final product viscosity. One way to solve this problem is by keeping the hydrolysis under control through the addition of a selected pH modifier.

